



umpectomy/ Reexcision

Instructions Before Surgery

The following information will help you to prepare for your upcoming surgery. We hope this information may help to make this experience more comfortable for you. If you have any questions regarding the instructions, please contact the Breast Care Center staff.

- This procedure is a same day surgery procedure. You may stay longer if it is medically necessary. You will need to have a responsible adult with you to drive you home. It is unsafe and against hospital policy to permit you to drive home after surgery when you have received any medication that might slow your responses (anesthesia, pain medication, or any medication to relieve anxiety).
- **NOTHING TO EAT OR DRINK AFTER MID-NIGHT** on the night before your surgery. Any medications that you take routinely should be taken at the usual time with a sip or two of water. People with diabetes, heart disease, and other illnesses should contact their primary care physicians for directions. Inform your physician if you are taking Coumadin® or other blood thinning medication.
- **DO NOT TAKE ASPIRIN** or aspirin containing products for 10 days before your surgery. Tylenol® is okay. STOP vitamin E supplements 2 weeks before or as soon as possible. Vitamin E in a multivitamin is okay.
- Wear comfortable clothing: a two-piece, loose outfit with a zipper or buttons in front is the easiest to put on. Please bring it with you.
- Recommended items to bring with you:
 - **DO NOT BRING VALUABLES, or give them to family/friends**
 - Tape player and headphones, favorite music, books on tape
 - Sweater or sweatshirt for warmth
 - Light reading
 - Important telephone numbers
 - A supportive bra **NO UNDERWIRE**, (sports bra or bra that hooks in front) **PLEASE WEAR A BRA FOR 3-4 DAYS FOLLOWING SURGERY, EVEN WHILE YOU SLEEP.** This minimizes post-operative bleeding and will make you more comfortable.

Lumpectomy/ Partial Mastectomy/ Reexcision

Instructions After Surgery

Pain Management

- People experience different types and amount of pain or discomfort after surgery. The goal of pain management is to assess your own level of discomfort and to take medication as it is needed. You will have better results controlling your pain if you take pain medication before your pain is severe.
- When pain is experienced, it is recommended to take pain medication on a regular schedule. Ibuprofen (e.g. Motrin®, Advil®) or Acetaminophen (e.g. Tylenol®) are non narcotics that can decrease pain. You will also be given a prescription for Vicodin® or another narcotic for the management of moderate pain.
- An ice pack may be helpful to decrease discomfort and swelling.
- Please notify us of any drug allergies, reactions or medical problems that would prevent you from taking these drugs. Vicodin® is a narcotic and should not be taken with alcoholic drinks. Do not use narcotics while driving.
- Narcotics can also cause or worsen constipation, so try to increase your fluids, eat high fiber foods (like prunes and bran), and make sure you are up, out of bed, taking small walks.
- Everyone is different and if one plan to decrease your pain is not working, it will be changed. Healing and recovery are improved with good pain control.

Incision (Scar) and Dressing Care

- Your incision (scar) has both stitches and steri-strips (small white strips of tape) and is covered by a gauze dressing and either a plastic dressing or tape.
- **Gauze and plastic dressing:** DO NOT REMOVE THE DRESSING, STERI STRIPS OR STITCHES. If the dressing falls off on its own, do not attempt to replace it.
- **Gauze and paper tape dressing:** You may remove the dressing 2 days after surgery. BE CAREFUL NOT TO TOUCH OR REMOVE THE STERI-STRIPS OR SUTURES.
- You may shower two days after surgery. Pat dry your plastic dressing or steri strips.
- Bruising and some swelling are common in women after surgery.
- A low-grade fever (under 100°F) is normal the day after surgery.

Activity

- Avoid strenuous activity, heavy lifting, and vigorous exercise until the stitches are removed. Tell us what you do and we will help you make a personal plan for “what you can do when” after surgery.
- Walking is a normal activity that can be restarted right away.
- No strenuous housework until after your first post-operative visit. No driving while you are on medication.
- Most people return to work within 1-2 weeks. Return to work varies with your type of work, your overall health and personal preferences. Discuss returning to work with us.

Diet

- You may resume your regular diet as soon as you can take fluids after recovering from anesthesia.
- We encourage 8-10 glasses of water and non-caffeinated beverages per day, plenty of fruits and vegetables, and lower fat foods. Talk with us about recommendations for healthy eating.
- A nutritionist is available for consultation in the Breast Care Center. Call the front desk to schedule an appointment.

Follow-Up Care

- The pathology results from your surgery should be available within **one week** after your surgery.
- We will contact you by telephone with the results or will inform you at your post-operative visit. Please let us know the telephone number where you may be reached with the results.
- Follow-up appointments are generally made before surgery with your physician and a nurse. Your sutures will be removed in approximately 10-14 days. Call the Breast Care Center if you do not have or remember an appointment.

Call the surgery number at the Breast Care Center for the following problems, any unanswered questions, or emotional support needs. Ask to speak with a nurse during the day. After business hours, your call will be forwarded to the answering service to reach your doctor or the doctor on call.

- **Pain that is not relieved by medication**
- **Fever more than 100° F or chills**
- **Excessive bleeding (a bloody dressing)**
- **Excessive swelling**
- **Redness outside the dressing**
- **Discharge or bad odor from the wound**
- **Allergic or other reactions to medication(s)**
- **Constipation (no bowel movements)**
- **Anxiety, depression, trouble sleeping, need more support**

